


CLAIMS

What is claimed is:

1. A combination, comprising valdecoxib, or a pharmaceutically acceptable salt thereof, and an allosteric carboxylic inhibitor of MMP-13 of Formula IC



or a pharmaceutically acceptable salt thereof, or an N-oxide thereof,
in which:

- 10 **R₁** represents a group selected from :
 - hydrogen, amino,
 - (C₁-C₆)alkyl, (C₃-C₆)alkenyl, (C₃-C₆)alkynyl, mono(C₁-C₆)alkylamino(C₁-C₆)alkyl, di(C₁-C₆)alkylamino(C₁-C₆)alkyl, aryl, aryl(C₁-C₆)alkyl, heterocycle, and 3- to 6-membered cycloalkyl(C₁-C₆)alkyl, these groups being unsubstituted or
- 15 substituted with one or more groups, which may be identical or different, selected from amino, (C₁-C₆)alkyl, cyano, halo(C₁-C₆)alkyl, C(=O)OR₄, OR₄ and SR₄, in which R₄ represents hydrogen or (C₁-C₆)alkyl,

W represents an oxygen atom, a sulphur atom, or a group =N-R', in which R' represents (C₁-C₆)alkyl, hydroxyl, or cyano,

- 20 **X₁, X₂ and X₃** represent, independently of each other, a nitrogen atom or a group -C-R₆ in which R₆ represents a group selected from hydrogen, (C₁-C₆)alkyl, amino, mono(C₁-C₆)alkylamino, di(C₁-C₆)alkylamino, hydroxyl, (C₁-C₆)alkoxy, and halogen,
- with the proviso that not more than two of the groups X₁, X₂ and X₃
- 25 simultaneously represent a nitrogen atom,

Y represents a group selected from oxygen atom, sulphur atom, -NH, and -N(C₁-C₆)alkyl,

Z represents:

- an oxygen atom, a sulphur atom,
- 5 • or a group -NR₇ in which R₇ represents a group selected from hydrogen, (C₁-C₆)alkyl, aryl(C₁-C₆)alkyl, cycloalkyl, aryl, and heteroaryl, and
- when Y is an oxygen atom, a sulphur atom, or a group -N(C₁-C₆)alkyl, Z optionally represents a carbon atom which is unsubstituted or substituted with a (C₁-C₆)alkyl, an aryl, an aryl(C₁-C₆)alkyl, an aromatic or non-aromatic
- 10 heterocycle or a cycloalkyl,

n is an integer from 1 to 8 inclusive,

Z₁ represents -CR₈R₉ wherein R₈ and R₉, independently of each other, represent a group selected from hydrogen, (C₁-C₆)alkyl, halo(C₁-C₆)alkyl, halogen, amino, OR₄, SR₄ or C(=O)OR₄ in which R₄ represents a hydrogen or (C₁-C₆)alkyl, and

- 15 • when n is greater than or equal to 2, the hydrocarbon chain Z₁ optionally contains one or more multiple bonds,
- and/or one of the carbon atoms in the hydrocarbon chain Z₁ may be replaced with an oxygen atom, a sulphur atom which is unsubstituted or substituted with one or two oxygen atoms, or a nitrogen atom which is
- 20 unsubstituted or substituted with a (C₁-C₆)alkyl,
- and when one of the carbon atoms in the hydrocarbon chain Z₁ is replaced with a sulphur atom which is unsubstituted or substituted with one or two oxygen atoms, then the group -C(=Y)-Z- optionally may be absent in the general formula (I),

25 A represents a group selected from :

- aromatic or non-aromatic, 5- or 6-membered monocycle comprising from 0 to 4 heteroatoms selected from nitrogen, oxygen and sulphur, and


- bicycle, composed of two aromatic or non-aromatic, 5- or 6-membered rings, which may be identical or different, comprising from 0 to 4 heteroatoms selected from nitrogen, oxygen and sulphur,

m is an integer from 0 to 7 inclusive,

- 5 **the group(s) R_2** , which may be identical or different, is (are) selected from (C₁-C₆)alkyl, halogen, -CN, NO₂, SCF₃, -CF₃, -OCF₃, -NR₁₀R₁₁, -OR₁₀, -SR₁₀, -SOR₁₀, -SO₂R₁₀, -(CH₂)_kSO₂NR₁₀R₁₁, -X₅(CH₂)_kC(=O)OR₁₀, -(CH₂)_kC(=O)OR₁₀, -X₅(CH₂)_kC(=O)NR₁₀R₁₁, -(CH₂)_kC(=O)NR₁₀R₁₁, and -X₄-R₁₂ in which:
- 10 • X₅ represents a group selected from oxygen, sulphur optionally substituted by one or two oxygen atoms, and nitrogen substituted by hydrogen or (C₁-C₆)alkyl,
- k is an integer from 0 to 3 inclusive,
 - R₁₀ and R₁₁, which may be identical or different, are selected from
- 15 hydrogen and (C₁-C₆)alkyl,
- X₄ represents a group selected from single bond, -CH₂-, oxygen atom, sulphur atom optionally substituted by one or two oxygen atoms, and nitrogen atom substituted by hydrogen atom or (C₁-C₆)alkyl group,
 - R₁₂ represents an aromatic or non-aromatic, heterocyclic or non-
- 20 heterocyclic, 5- or 6-membered ring which is unsubstituted or substituted with one or more groups, which may be identical or different, selected from (C₁-C₆)alkyl, halogen, hydroxyl and amino, and when the ring is heterocyclic, it comprises from 1 to 4 heteroatoms selected from nitrogen, oxygen and sulphur;

R₃ represents a group selected from:

- hydrogen,
 - (C₁-C₆)alkyl, (C₃-C₆)alkenyl, (C₃-C₆)alkynyl, these groups being unsubstituted or substituted with one or more groups, which may be identical or different, selected from amino, cyano, halo(C₁-C₆)alkyl, cycloalkyl, -
- 5 C(=O)NR₁₀R₁₁, -C(=O)OR₁₀, OR₁₀, and SR₁₀, in which R₁₀ and R₁₁, which may be identical or different, represent hydrogen or (C₁-C₆)alkyl,
- and the group of formula :



- ✓ in which p is an integer from 0 to 8 inclusive,
- 10 ✓ Z₂ represents -CR₁₃R₁₄ wherein R₁₃ and R₁₄, independently of each other, represent a group selected from hydrogen, (C₁-C₆)alkyl, phenyl, halo(C₁-C₆)alkyl, halogen, amino, OR₄, SR₄ and -C(=O)OR₄ in which R₄ represents hydrogen or (C₁-C₆)alkyl, and
- when p is greater than or equal to 2, the hydrocarbon chain Z₂ optionally
- 15 contains one or more multiple bonds,
- and/or one of the carbon atoms in the hydrocarbon chain Z₂ may be replaced with an oxygen atom, a sulphur atom which is unsubstituted or substituted with one or two oxygen atoms, a nitrogen atom which is unsubstituted or substituted with a (C₁-C₆)alkyl, or a carbonyl group,
- 20 ✓ B represents a group selected from:
- an aromatic or non-aromatic 5- or 6-membered monocycle comprising from 0 to 4 heteroatoms selected from nitrogen, oxygen and sulphur, and
 - a bicycle, composed of two aromatic or non-aromatic, 5- or 6-membered rings, which may be identical or different, comprising from 0 to 4
- 25 heteroatoms selected from nitrogen, oxygen and sulphur,
- ✓ q is an integer from 0 to 7 inclusive,

- ✓ the group(s) R_5 , which may be identical or different, is (are) selected from (C_1-C_6) alkyl, halogen, CN, NO_2 , CF_3 , OCF_3 , $-(CH_2)_kNR_{15}R_{16}$, -
 $N(R_{15})C(=O)R_{16}$, $-N(R_{15})C(=O)OR_{16}$, $-N(R_{15})SO_2R_{16}$, $-N(SO_2R_{15})_2$, -
 OR_{15} , $-S(O)_{k1}R_{15}$, $-SO_2-N(R_{15})-(CH_2)_{k2}-NR_{16}R_{17}$, -
5 $(CH_2)_kSO_2NR_{15}R_{16}$, $-X_7(CH_2)_kC(=O)OR_{15}$, -
 $(CH_2)_kC(=O)OR_{15}$, $-C(=O)O-(CH_2)_{k2}-NR_{15}R_{16}$, $-C(=O)O-(CH_2)_{k2}-C(=O)OR_{18}$,
 $-X_7(CH_2)_kC(=O)NR_{15}R_{16}$, $-(CH_2)_kC(=O)NR_{15}R_{16}$, $-R_{19}-C(=O)OR_{15}$, $-X_6-R_{20}$,
and $-C(=O)-R_{21}-NR_{15}R_{16}$ in which :
- 10 - X_7 represents a group selected from oxygen atom, sulphur atom
optionally substituted by one or two oxygen atoms, and nitrogen atom
substituted by a hydrogen atom or a (C_1-C_6) alkyl group,
- k is an integer from 0 to 3 inclusive,
- $k1$ is an integer from 0 to 2 inclusive,
- $k2$ is an integer from 1 to 4 inclusive,
- 15 - R_{15} , R_{16} and R_{17} , which may be identical or different, are selected from
hydrogen and (C_1-C_6) alkyl,
- R_{18} represents a group selected from (C_1-C_6) alkyl, $-R_{21}-NR_{15}R_{16}$,
 $-R_{21}-NR_{15}-C(=O)-R_{21}-NR_{16}R_{17}$, and $-C(=O)O-R_{21}-NR_{15}R_{16}$ in which R_{21}
represents a linear or branched (C_1-C_6) alkylene group, and R_{15} , R_{16} and R_{17}
20 are as defined hereinbefore,
- R_{19} represents a (C_3-C_6) cycloalkyl group,
- X_6 represents a group selected from single bond, $-CH_2-$, oxygen atom,
sulphur atom optionally substituted by one or two oxygen atoms, and
nitrogen atom substituted by hydrogen atom or (C_1-C_6) alkyl group,

- R₂₀ represents an aromatic or non-aromatic, heterocyclic or non-heterocyclic, 5- or 6-membered ring, which is unsubstituted or substituted with one or more groups, which may be identical or different, selected from (C₁-C₆)alkyl, halogen, hydroxyl, oxo, cyano, tetrazole, amino, and -
5 C(=O)OR₄ wherein R₄ represents hydrogen or (C₁-C₆)alkyl, and, when the ring is heterocyclic, it comprises from 1 to 4 heteroatoms selected from nitrogen, oxygen and sulphur,

with the proviso that when X₁ represents a nitrogen atom, X₂ cannot represent a carbon atom substituted with a methyl group or with NH-CH₃.

10

2. The combination according to Claim 1, wherein the compound of Formula IC is selected from:

4-[6-(4-Methoxy-benzylcarbamoyl)-1-methyl-2,4-dioxo-1,4-dihydro-2H-pyrido[3,4-d]pyrimidin-3-ylmethyl]-benzoic acid;

15 3-Benzyl-1-methyl-2,4-dioxo-1,2,3,4-tetrahydro-pyrido[3,4-d]pyrimidine-6-carboxylic acid (1,3-benzodioxol-5-ylmethyl)-amide;

Methyl 4-[6-(4-Methoxy-benzylcarbamoyl)-1-methyl-2,4-dioxo-1,4-dihydro-2H-pyrido[3,4-d]pyrimidin-3-ylmethyl]-benzoate;


20 3-(4-Cyano-benzyl)-1-methyl-2,4-dioxo-1,2,3,4-tetrahydro-pyrido[3,4-d]pyrimidine-6-carboxylic acid 4-methoxy-benzylamide;

4-[6-(3-Methoxy-benzylcarbamoyl)-1-methyl-2,4-dioxo-1,4-dihydro-2H-pyrido[3,4-d]pyrimidin-3-ylmethyl]-benzoic acid;

4-[6-(4-Methoxy-benzylcarbamoyl)-1-methyl-2,4-dioxo-1,4-dihydro-2H-pyrido[2,3-d]pyrimidin-3-ylmethyl]-benzoic acid;

25 or a pharmaceutically acceptable salt thereof.

3. A combination, comprising valdecoxib, or a pharmaceutically acceptable salt thereof, and an allosteric carboxylic inhibitor of MMP-13 of Formula VG



VG

or a pharmaceutically acceptable salt thereof, wherein

R^1 and R^2 independently are hydrogen, halo, hydroxy, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, NO_2 , NR^4R^5 , CN, or CF_3 ;

n is 1, and

Each Ar independently is aryl or Het, wherein aryl is phenyl or substituted phenyl, and Het is an unsubstituted or substituted heteroaryl group.

- 10 4. A pharmaceutical composition, comprising a combination of valdecoxib, or a pharmaceutically acceptable salt thereof, and an allosteric carboxylic inhibitor of MMP-13, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier, diluent, or excipient.
- 15 5. A method of treating a disease or disorder selected from cartilage damage, inflammation, arthritis, and pain in a mammal, comprising administering to the mammal a therapeutically effective amount of a combination of valdecoxib, or a pharmaceutically acceptable salt thereof, and an allosteric carboxylic inhibitor of MMP-13, or a pharmaceutically acceptable salt thereof.
- 20 6. The method according to Claim 5, wherein the disease or disorder is rheumatoid arthritis.
- 25 7. The method according to Claim 5, wherein the disease or disorder is osteoarthritis.

8. The method according to Claim 5, wherein the disease or disorder is joint inflammation.
9. The method according to Claim 5, wherein the pain is joint pain.